

## Harsukh Educational Charitable Society

### International Journal of Community Health and Medical Research

Journal home page: [www.ijchmr.com](http://www.ijchmr.com)

doi: 10.21276/ijchmr

ISSN E: 2457-0117 ISSN P:2581-5040

Index Copernicus ICV 2017=57.10

## Original Research

### Evaluation of stroke cases in emergency department- A clinical study

Jyoti Pankaj<sup>1</sup> Arun Kumar Bharti<sup>2</sup>

<sup>1</sup>Assistant Professor, Dept Of General Medicine, Prasad Institute Of Medical Sciences Lucknow <sup>2</sup>Junior Resident Dept Of Pharmacology KGMU Lucknow

#### ABSTRACT

**Background:** There is considerable increase in number of stroke cases. The present study was conducted to assess the cases of stroke reported to emergency department. **Materials & Methods:** The present study was conducted in the Emergency department on 112 cases of both genders. Parameters such as education, vascular risk factors and concomitant diseases (prior stroke/ TIA, smoking, heavy drinking, hypertension, coronary heart disease, peripheral vascular disease, diabetes mellitus, hyperlipidaemia etc. was recorded in cases file. **Results:** Out of 112 cases, males were 72 and females were 50. Maximum cases were seen in age group 60-80 years (68) followed by >80 years (24), 40-60 years (16) and 20-40 years (4). The difference was significant ( $P < 0.05$ ). Various risk factors were hypertension seen in 102, diabetes mellitus in 84, hyperlipidaemia in 92, drinking in 69, smoking in 58, low education status in 66 and previous stroke in 15 cases. The difference was significant ( $P < 0.05$ ). **Conclusion:** The number of stroke cases is increasing day by day. We found that various risk factors were hypertension, diabetes mellitus, hyperlipidaemia, drinking, smoking, low education status and previous stroke.

**Key words:** Emergency, Hypertension, Stroke

**Corresponding author:** Dr. Arun Kumar Bharti, Junior Resident Dept Of Pharmacology KGMU Lucknow

**This article may be cited as:** Pankaj J , Bharti AK. Evaluation of stroke cases in emergency department- A clinical study .HECS Int J Comm Health Med Res 2019; 5(2):1-3

#### INTRODUCTION

World Health Organization defined stroke as a "neurological deficit of cerebrovascular cause that persists beyond 24 hours or is interrupted by death within 24 hours", although the word "stroke" is centuries old. This definition was supposed to reflect the reversibility of tissue damage and was devised for the purpose, with the time frame of 24 hours being chosen arbitrarily.<sup>1</sup> The 24-hour limit divides stroke from transient ischemic attack, which is a related syndrome of stroke symptoms that resolve completely within 24 hours. With the availability of treatments which can reduce stroke severity when given early, many now prefer alternative terminology, such as brain attack and acute ischemic cerebrovascular syndrome (modeled after heart attack and acute coronary syndrome, respectively), to reflect the urgency of stroke symptoms and the need to act swiftly.<sup>2</sup> Strokes can be classified into two major categories: ischemic and hemorrhagic. Ischemic strokes are caused by interruption of the blood supply to the brain, while hemorrhagic strokes result from the rupture of a blood vessel or an abnormal vascular structure.

About 87% of strokes are ischemic, the rest being hemorrhagic. Bleeding can develop inside areas of ischemia, a condition known as "hemorrhagic transformation." It is unknown how many hemorrhagic strokes actually start as ischemic strokes.<sup>3</sup> The present study was conducted to assess the cases of stroke reported to emergency department.

#### MATERIALS & METHODS

The present study was conducted in the Emergency department. It comprised of 112 cases of both genders. Attendants were informed regarding the study and written consent was obtained. Ethical clearance was obtained prior to the study. General information such as name, age, gender etc. was recorded. Parameters such as education, vascular risk factors and concomitant diseases (prior stroke/ TIA, smoking, heavy drinking, hypertension, coronary heart disease, peripheral vascular disease, diabetes mellitus, hyperlipidaemia etc. was recorded in cases file. Results thus obtained were subjected to statistical analysis. P value less than 0.05 was considered significant ( $P < 0.05$ ).

**RESULTS**

**Table I Distribution of cases**

Total- 112		
Gender	Males	Females
Number	72	50

Table I shows that out of 112 cases, males were 72 and females were 50.

**Table II Age wise distribution of cases**

Age group (Years)	Number	P value
20-40	4	0.01
40-60	16	
60-80	68	
>80	24	


Table II shows that maximum cases were seen in age group 60-80 years (68) followed by >80 years (24), 40-60 years (16) and 20-40 years (4). The difference was significant (P< 0.05).

**Table III Assessment of risk factors**

Parameters	Number	P value
Education Higher	46	0.01
Primary	66	
Smoking	58	
Drinking	69	
Hypertension	102	
Diabetes mellitus	84	
Hyperlipidemia	92	
Previous stroke	15	

Table III, graph I shows that various risk factors were hypertension seen in 102, diabetes mellitus in 84, hyperlipidaemia in 92, drinking in 69, smoking in 58, low education status in 66 and previous stroke in 15 cases. The difference was significant (P< 0.05).

**Graph I Risk factors**



**DISCUSSION**

Stroke is the second leading cause of death worldwide. In last few years, the incidence of stroke and burden of stroke increased in both developing and developed countries. Improving quality of care is critical to improve the prognosis after stroke. Several studies of patients with stroke have reported higher mortality on weekends and weekday nights. A recent systematic review and meta-analysis suggests that patients with acute ischemic stroke presenting during off-hours had higher short-term mortality and disability compared with patients presenting during regular hours.<sup>4</sup> The result was robust across subgroups and sensitivity analyses. A higher incidence of mortality and a decreased use of invasive cardiac procedures was noticed in patients admitted with myocardial infarction during weekends, a phenomenon termed the “weekend effect”.<sup>5</sup> The present study was conducted to assess the cases of stroke reported to emergency department.

We observed that out of 112 cases, males were 72 and females were 50. Maximum cases were seen in age group 60-80 years (68) followed by >80 years (24), 40-60 years (16) and 20-40 years (4). Various risk factors were hypertension seen in 102, diabetes mellitus in 84, hyperlipidaemia in 92, drinking in 69, smoking in 58, low education status in 66 and previous stroke in 15 cases.

Albright et al<sup>6</sup> reported that epileptic seizures (41%) and psychiatric disorders (28%) are the most common diseases that can cause stroke mimics, and the frequency of hypoglycemia is low (1%).

Palmer et al<sup>7</sup> found that off-hour presentation was identified in 2672 (59.5%) patients with ischemic stroke. Comparison of patients admitted during off-hours with those admitted during on-hours revealed an unadjusted odds ratio of in-hospital mortality of 1.38, which declined to 1.34 after adjusting for patient characteristics. No difference in 30-day mortality, total death or dependence at three, six and 12 months between two groups was observed. No association between off-hour admission and quality of care was found.

Martinez<sup>8</sup> observed that of 1,557 suspected acute stroke cases examined at the emergency department, 137 (8.8%) were stroke mimics. The most common causes were symptomatic epilepsy (28 cases, 20.4%), neuropathy-like symptoms (21 cases, 15.3%), and hypoglycemia (15 cases, 10.9%). Outcomes were survival to hospital discharge for 91.2% and death for 8.8% of the cases. Clinical results were significantly different between stroke mimics and the stroke group for low systolic blood pressure, low National Institutes of Health Stroke Scale score on initial treatment, history of diabetes, and no history of arrhythmia. On multivariate analysis, distinguishing factors for stroke mimics include systolic blood pressure ≤ 140 mmHg, National Institutes of Health Stroke Scale score ≤ 5 points, history of diabetes, and no history of arrhythmia.

**CONCLUSION**

The number of stroke cases is increasing day by day. We found that various risk factors were hypertension, diabetes mellitus, hyperlipidaemia, drinking, smoking, low education status and previous stroke.

**REFERENCES**

1. Recommendations on stroke prevention, diagnosis, and therapy. Report of the WHO Task Force on Stroke and other Cerebrovascular Disorders. Stroke. 1989; 20: 1407–1431.

2. Saposnik G, Baibergenova A, Bayer N, Hachinski V. Weekends: A dangerous time for having a stroke? *Stroke*. 2007; 38: 1211–1215.
3. Reeves MJ, Smith E, Fonarow G, Hernandez A, Pan W, Schwamm LH, et al. Off-hour admission and in-hospital stroke case fatality in the get with the guidelines-stroke program. *Stroke*. 2009; 40: 569–576.
4. Jauss M, Oertel W, Allendoerfer J, Misselwitz B, Hamer H. Bias in request for medical care and impact on outcome during office and non-office hours in stroke patients. *Eur J Neurol* 2009; 16: 1165–1167.
5. McKinney JS, Deng Y, Kasner SE, Kostis JB, Myocardial Infarction Data Acquisition System Study G. Comprehensive stroke centers overcome the weekend versus weekday gap in stroke treatment and mortality. *Stroke* 2011; 42: 2403–2409.
6. Albright KC, Raman R, Ernstrom K, Hallevi H, Martin-Schild S, Meyer DM, et al. Can comprehensive stroke centers erase the 'weekend effect'? *Cerebrovasc Dis*. 2009; 27: 107–113.
7. Palmer WL, Bottle A, Davie C, Vincent CA, Aylin P. Dying for the weekend: a retrospective cohort study on the association between day of hospital presentation and the quality and safety of stroke care. *Arch Neurol*. 2012; 69: 1296–1302.
8. Martinez-Martinez MM, Fernandez-Travieso J, Fuentes B, Ruiz-Ares G, Martinez-Sanchez P, Cazorla Garcia R, et al. Off-hour effects on stroke care and outcome in stroke centres. *Eur J Neurol*. 2012; 19:1140–1145.