

Harsukh Educational Charitable Society

International Journal of Community Health and Medical Research

Journal home page: www.ijchmr.com

doi: 10.21276/ijchmr

ISSN E: 2457-0117 ISSN P: 2581-5040

Index Copernicus ICV 2018=62.61

Original ARTICLE

A STUDY ON THE PREVALENCE OF PSYCHIATRIC COMORBIDITIES AMONG CANNABIS USERS

Rakesh Kumar¹, Ripu Daman Singh Dhariwal², Kamal Kumar Verma³, Preetam Singh⁴, Jitendra Acharya⁵

¹Assistant professor, Department of Psychiatry, S P Medical College Bikaner, Rajasthan, India;

²Junior Specialist, Department of Psychiatry, District hospital, Shri Ganganagar, Rajasthan, India;

³Senior professor, Department of Psychiatry, S P Medical College Bikaner, , Rajasthan, India;

⁴Resident, Department of Psychiatry, S P Medical College Bikaner, , Rajasthan, India;

⁵Department of Dentistry, S.P. Medical College Bikaner, Rajasthan, India

ABSTRACT

Introduction: The prevalence of psychiatric disorder beside with co-occurring substance use disorders and vice-versa are growing through time which worsen treatment outcome and boost relapse. **Material And Method:** Total 40 cases of cannabis addiction who attended psychiatry opd of S.P. Medical College Bikaner were selected based on ICD 10 criteria. The psychiatric comorbidities were assessed in them using MINI INTERNATIONAL NEUROPSYCHIATRIC INTERVIEW. **Result:** In this study of 40 cannabis dependent patients psychiatric comorbidities were found in 39(97.5%). Over all comorbidities were mood disorder in 18 (45%), psychotic disorder 16 (40%), antisocial personality disorder 3 (7.5%) and anxiety disorder in 2 (5%). **Conclusion:** From our study we came to the conclusion that psychiatric comorbidities are common among cannabis dependent patients.

KEYWORDS: cannabis dependent, psychiatric comorbidities.

Corresponding author Dr. Ripu Daman Singh Dhariwal, Junior Specialist, Department of Psychiatry, District hospital, Shri Ganganagar, Rajasthan, India

This article may be cited as: Kumar R, Dhariwal Rds, Verma Kk, Singh P, Acharya J. A Study On The Prevalence Of Psychiatric Comorbidities Among Cannabis Users. HECS Int J Comm Health Med Res 2019; 5(3):143-145.

INTRODUCTION

Comorbidity shows the presences of a diverse clinical entity that has existed or may occur during the clinical outcome of a patient with the index disease.¹ Psychiatric comorbidities are worldwide among substance users and vice-versa. There is specifics that taking cannabis leads to acute adverse mental effects in a high proportion of regular users.² In a study done on 44 schools in Victoria, Australia on 1601 students aged 14-15 years found 66% of males and 52% of females use cannabis with 10% of males and 4% of females using it daily. The prevalence of depression and anxiety was found to be more in females compared to males.³ It was also found that adolescent females are particularly at risk of developing depression from cannabis use though they may not use as heavily as their male peers.⁴ In National Comorbidity Survey (NCS) it was found that among those with a 12 month diagnosis of any substance use disorder, 36% had at least one anxiety disorder whilst 25% had at least one affective disorder.⁵ The aim of the study is to assess the

prevalence of psychiatric comorbidities among cannabis dependent patients.

MATERIALS AND METHOD

This is a descriptive and analytical cross-sectional study conducted on 40 patients who attended the psychiatry OPD of S.P. Medical College and Hospital, Bikaner from August 2016 to July 2017 and fulfilled the criteria for cannabis dependence based on ICD 10. The psychiatric comorbidities were assessed in them using MINI INTERNATIONAL NEUROPSYCHIATRIC INTERVIEW (7.0.0 for DSM-5). Semi-structured Performa was used for socio-demographic profile.

RESULTS


Out of 40 cannabis dependent patients, majority belonged to 17-49 years age group with mean age of 26.95 years. Of them all

were males, 82.5% were Hindus, and most of them were unmarried (77.5%). Table 1 has shown the socio-demographic profile of the patients. Out of 40 cannabis dependent patients, 39(97.5%) were having psychiatric comorbidities. Figure 1 has shown the prevalence of psychiatric comorbidities among cannabis dependent patients.

Table 1. Demographic Distribution

Category	Variables
Sex	Male- 40 Female- 0
Religion	Hindu- 33 Muslim- 7
Marital status	Married- 9, Unmarried- 31
Educational status	Illiterate- 15, Class<=12- 16, Class> 12- 9
Occupation	Unemployed 7, Employed- 28, Student- 5

FIGURE 1 Diagram showing prevalence of psychiatric comorbidities among cannabis dependent patients.



DISCUSSION

Our study sample mostly belonged to age group 17-49 years and all were males which is consistent with previous studies all reporting higher percentage of males in their studies.^{6,7}In the present study most of the patients were unmarried 31(77.5%) which is consistent with previous study which showed that cannabis use reduced with marriage.⁸ In the study, among the cannabis dependent patients, 15 (37.5%) were illiterate and 16 (40%) studied up to or less than class 12 which is consistent with studies reporting high dropout rates from school among adolescent cannabis users.^{9,10,11} In this study, among 40 cannabis dependent patients 18 (45%) have a comorbid mood disorder which include manic episode 3 (7.5%), bipolar type I with psychotic features 7 (17.5%), bipolar type I without psychotic features 5 (12.5%), bipolar type 2 (2.5%) and major depressive disorder 2 (5%) which is consistent with the findings of Guillem E et al. who conducted a study to see the prevalence of psychiatric disorders in 90 cannabis users and mood disorder was found in 48% of them.¹² In the present study, among 40 cannabis

dependent patients, 2 (5%) had comorbid generalized anxiety disorder whereas in the study done by Thomas et al. to see the comorbidities associated with cannabis use found that 22% had comorbid anxiety disorder.¹³ Antisocial personality disorder was found in 3(7.5%) whereas previous studies revealed that comorbid antisocial personality disorders were very common among substance dependent patients¹⁴. Psychotic disorder was found to be most prevalent among the cannabis dependent patients 16 (40%) which is consistent with previous study done by Le Bec et al. who reported that there was a relationship between cannabis and psychosis onset and cannabis consumption can also be an independent risk factor of psychotic disorders.¹⁵ In the study done by DeQuardo et al. found that among schizophrenia patients 20% of female and 48% of male patients had comorbid substance use and the most common substance was cannabis.¹⁶

CONCLUSION

We concluded that cannabis addiction is commonly associated with comorbid psychiatric disorder. So timely detection and treatment of psychiatric comorbidities are very necessary while treating the patients of cannabis dependence for the prevention of future relapses and for better treatment outcome. Better perceptiveness of how cannabis dependence disrupts some of the brain functions making an individual more prone to develop psychiatric illness and vice versa would help clinicians to offer better treatment to these patients. However, community based study is essential to conclude more correctly.

REFERENCES

1. Feinstein AR. The pre-therapeutic classification of comorbidity in chronic disease. *J Chronic Dis* 1970; 23 : 455.
2. Johns A. Psychiatric effects of cannabis. *British Journal of Psychiatry* 2001;178:116- 122.
3. Patton GC , Coffey C, Carlin J B , Degenhardt L, Lynskey M , Hall W. Cannabis use and mental health in young people :cohort study. *BMJ* 2002Nov 23;325(7374):1195-8.
4. McGee R, Williams S A, Poulton R. & Moffitt T. A longitudinal study of cannabis use and mental health from adolescence to early adulthood. 2000. *Addiction*;95: 491–503.
5. Kessler R.C, McGonagle K.A., Zhao S, Nelson C.B., Hughes M, Eshleman S,Wittchen H.U, Kendler K.S. Lifetime and 12-month prevalence of DSM-III-R psychiatric disorders in the United States. Results from the National Comorbidity Survey. *Archives of General Psychiatry*, 1994 51(1), 8–19.
6. Chaturvedi HK, Phukan RK, Mahanta J. The association of selected sociodemographic factors and differences in patterns of substance use: a pilot study in selected areas of Northeast India. *Substance Use and Misuse*. 2003; 38(9):1305–1322.
7. Patton GC , Coffey C, Carlin J B , Degenhardt L, Lynskey M , Hall W. Cannabis use and mental health in young people :cohort study. *BMJ* 23 NOVEMBER 2002;325.
8. Duncan GJ, Wilkerson B, England P. Cleaning up their act: the effects of marriage and cohabitation on licit and illicit drug use. *Demography* 2006;43:691–710.
9. Brook JS, Balka EB, Whiteman M. The risks for late adolescence of early marijuana use. *American Journal of Public Health* 1999; 89: 1549-1554.
10. Bray J, Zarkin G, Ringwalt C, Qi J. The relationship between marijuana initiation and dropping out of high school. *Health Economics* 2000; 9: 9-18.

11. Yamada T, Kendix M. The impact of alcohol and marijuana consumption on high school graduation. *Health Economics*, 1998.
12. Guillem E, Pelissolo A, Vorspan F, Bouchez-Arbabzadeh S, Lépine JP. Sociodemographic profiles, addictive and mental comorbidity in cannabis users in an outpatient specific setting. *Encephale*. 2009 Jun;35(3):226-33.
13. Thomas H. Psychiatric symptoms in cannabis users. 1993. *British journal of psychiatry*,163, 141-149.
14. Compton WM Cottler LB, Phelps DL, Ben Abdallah A, Spitznagel EL. Psychiatric disorders among drug dependent subjects: are they primary or secondary? *Am J Addict*. 2000 Spring;9(2):126-34.
15. Le Bec PY, Fatseas M, Denis C, Lavie E, Auriacombe M. Cannabis and psychosis: search of a causal link through a critical and systematic review. *Encephale*. 2009;35(4):377–85.
16. DeQuardo JR, Carpenter CF, Tandon R. Patterns of substance abuse in schizophrenia: nature and significance. *J Psychiatr Res*. 1994;28(3):267–75.