

Harsukh Educational Charitable Society

International Journal of Community Health and Medical Research

Journal home page: www.ijchmr.com

doi: 10.21276/ijchmr

ISSN E: 2457-0117 ISSN P: 2581-5040

Index Copernicus ICV 2017=57.10

Original Research

Evaluation of risk factors of stroke - A clinical study

Deepak K. Gautam¹, Parul Sharma², Pankaj K. Mishra³

¹Associate Professor, Dept. of General Medicine, ²Assistant Professor, Dept. of Physiology, Institute of Medical Sciences, Banaras Hindu University, Varanasi, U.P., India

³Professor, Dept. of Community Medicine, Mayo Institute of Medical Sciences, Faizabad Road, Barabanki, Lucknow, U.P., India

ABSTRACT

Background: Ischemic stroke is the most common type of stroke, accounting for more than three-fourth of all cases. The present study was conducted to assess the cases of stroke. **Materials & Methods:** The present study was conducted on 122 cases of strokes of both genders. Sociodemographic profile, individual vascular risk factors, medical history and clinical assessments were studied. **Results:** Out of 122 patients, males were 82 and females were 40. TIA was seen in 28 males and 12 females, ischemia in 40 males and 8 females and hemorrhage in 14 males and 10 females. The difference was significant ($P < 0.05$). Common risk factors were previous stroke seen in 24, previous TIA in 12, IHD in 5, atrial fibrillation in 8, peripheral artery disease in 7, migraine in 4, diabetes mellitus in 6, hypertension in 36 and smoking in 20. **Conclusion:** Stroke is a leading cause of morbidity and mortality. Common types are TIA, ischemic and hemorrhagic.

Key words: hemorrhagic, ischemic, stroke

Corresponding author: Dr. Parul Sharma, Assistant Professor, Dept. of Physiology, Institute of Medical Sciences, Banaras Hindu University, Varanasi

This article may be cited as: Gautam DK, Sharma P, Mishra PK. Evaluation of risk factors of stroke - A clinical study. HECS Int J Comm Health Med Res 2019; 5(1):22-25

INTRODUCTION

Stroke is a heterogeneous group of cerebrovascular diseases with a wide variety of phenotypes and underlying etiologies. Ischemic stroke is the most common type, representing more than 80% of all stroke cases. It has been defined as rapidly developing signs of focal or global disturbance of cerebral function with symptoms lasting for ≥ 24 hours, or leading to death with apparently no cause other than that of vascular origin.¹

Factors such as lifestyles, associated risk factors, differences in study designs and stroke classification tools have shown varied associations between the risk factors of stroke and ischemic subtypes. However, hypertension and diabetes mellitus were linked more to small vessel disease, whereas large artery atherosclerosis had a predilection for being associated with hyperlipidemia and smoking.²

It is a collection of clinical syndromes resulting from cerebral ischemia to intracranial hemorrhage. It is the 3rd most common cause of morbidity and mortality in the west. Some of the studies done recently have explicated the pattern of stroke to a considerable extent in our country with a prevalence rate of 471.58 per lakh population.³

Common symptoms of Acute Ischemic Stroke include hemiparesis, monoparesis or quadriplegia, hemisensory deficits, monocular or binocular visual loss, visual field defects, diplopia, dysarthria, facial droop, ataxia, vertigo, aphasia and sudden decrease in the level of consciousness. Conditions that may mimic stroke are Bell's palsy, conversion disorder, complicated migraine, psychogenic conditions, hypoglycemia, CNS infections or brain abscess, hypertensive encephalopathy, seizures or tumor.⁴ The present study was undertaken to assess the stroke cases admitted to

the Casualty and General Medicine department of a tertiary care setup.

MATERIALS & METHODS

The present study was conducted in the department of Community Medicine, Mayo Institute of Medical Sciences, Lucknow. It comprised of 122 cases of stroke of both genders. Ethical clearance of the institute’s ethical committee was taken prior to starting the study. Information pertaining to particulars of the subjects such as name, age, gender etc. were recorded. Sociodemographic characteristics, individual vascular risk factors, medical history and clinical assessments were done. Patients of ischemic stroke were divided into subgroups. A stroke was considered ischemic if brain imaging revealed acute infarction or did not reveal any evidence of hemorrhage. Results obtained were subjected further to statistical analysis. P value of <0.05 was considered to be significant.

Total - 122		
Gender	Males	Females
Number	82	40

Table I shows that out of 122 patients, males were 82 and females were 40.

Table II: Type of stroke


Type	Male	Female	P value
TIA	28	12	0.01
Ischemic	40	8	
Hemorrhagic	14	10	
Total	82	40	

RESULTS


Table I: Distribution of patients

Table II, graph I shows that TIA was seen in 28 males and 12 females, ischemic in 40 males and 8 females and hemorrhagic in 14 males and 10 females. The difference was significant (P< 0.05).

Graph I: Type of stroke



Graph II: Risk factors in patients



Graph II shows that common risk factors were previous stroke seen in 24, previous TIA in 12, IHD in 5, atrial fibrillation in 8, peripheral artery disease in 7, migraine in 4, diabetes mellitus in 6, hypertension in 36 and smoking in 20.

DISCUSSION

According to the definition proposed by the World Health Organization in 1970, “stroke is rapidly developing clinical signs of focal (or global) disturbance of cerebral function, with symptoms lasting 24 hour or longer, or leading to death, with no apparent cause other than of vascular origin.” Recently, a new definition of stroke has been proposed by the American Stroke Association which is tissue based and states that “stroke is an episode of acute neurological dysfunction presumed to be caused by ischemia or hemorrhage, persisting for ≥ 24 hours or until death.” The clinical manifestations of stroke include a wide range of signs and symptoms such as paralysis, weakness, numbness, severe headache, slurred speech, mental status change, vision problems, falling and dizziness⁵. A stroke may be described as silent if it occurs without clinical features. In silent strokes, either a silent area of the brain is involved or symptoms remain inconspicuous. The present study was undertaken to assess the stroke cases admitted to the General Medicine department of a tertiary care center. We found that out of 122 patients, males were 82 and females were 40. In a recent study conducted in India, China and Latin-America, chronic diseases (stroke, heart disease, diabetes mellitus, chronic respiratory disease and malignancy) as a whole accounted for the majority of death among the elderly⁵. Individual Indian studies have estimated that the prevalence rates increase from 0.1-0.3 per thousand in the under 45-year age group to 12-20 per thousand in 75-84 year age group. In India, stroke in younger people is high (18-32%) of all stroke cases, compared to high income countries. Indian men are more likely to have a stroke than their women counterpart: the male/female sex ratio for India is 7:1.8. It can be explained to be due to the differences in risk factors such as drinking and smoking which are more prevalent among men than women in our country⁶. Chitrabalam et al⁷

found that TIA was seen in 28 males and 12 females, ischemic in 40 males and 8 females and hemorrhagic in 14 males and 10 females. Common risk factors were previous stroke seen in 24, previous TIA in 12, IHD in 5, atrial fibrillation in 8, peripheral artery disease in 7, migraine in 4, diabetes mellitus in 6, hypertension in 36 and smoking in 20. Baijuet et al⁸ found that incidence of stroke in elderly (≥ 45 years) was than young (< 45 years) patients with male predominance in both the groups. The most common presentation from both age groups was hemiplegia (75% in young vs. 72.1% in elderly). Common risk factors were hypertension and past history of CVD (25%) and smoking (16.6%) in young and hypertension (37.1%), dyslipidemia and smoking (14.3%) in elderly. The most common type of stroke in both age groups was ischemic (83.3% in young vs. 73.2% in elderly), followed by hemorrhagic. Common types of hemorrhage in both the young as well as the elderly on the basis of site of bleed were lobar and thalamoganglionic. The most common site of infarction in both age group is MCA territory. Intracerebral hemorrhage is mostly caused by uncontrolled hypertension leading to rupture of small vessels. An ‘avalanche type effect’ results due to this rupture with breakage of nearby vessels resulting in expansion of hematoma in up to 40% of cases. Subarachnoid hemorrhage is mainly due to saccular aneurysms though it is also associated with arteriovenous malformation, intracranial neoplasm and some medications such as anticoagulants. About 65% of subarachnoid hemorrhage patients survive, but half remain disabled primarily due to severe cognitive deficit.⁹

CONCLUSION

Stroke is a leading cause of morbidity and mortality. Common types are TIA, ischemic and hemorrhagic.

REFERENCES

1. S. K. Das, T. K. Banerjee, A. Biswas, D. K. Raut, C. S. Mukherjee, A. Chaudhari, et al. A prospective community based study of stroke in Kolkata, India. Stroke. 2007; 38(2):906-10.

2. D. Nagaraja, G. Gururaj, N. Girish, Samhita Panda, A. K. Roy, G. R. K. Sarma, et al. Feasibility study of stroke surveillance: data from Bangalore, India. *Indian J Med Res.* 2009 Oct; 130:396-403.
3. Ferri CP, Acosta D, Guerra M, Huang Y, Llibre-Rodriguez JJ, Salas A, et al. Socioeconomic factors and all cause and cause-specific mortality among older people in Latin America, India, and china: a population-based cohort study. *PLoS Med.* 2012; 9(2):1001179.
4. Prasad Kameshwar, Singhal Kapil K. Stroke in young: an Indian perspective. *Neurol India.* 2010; 58(3):343-50.
5. Dalal PM, Malik S, Bhattacharjee M, Trivedi ND, Vairale J, Bhat P, et al. Population-based stroke survey in Mumbai, India: incidence and 28-day case fatality. *Neuroepidemiology.* 2008; 31:254-61.
6. Sethi P. Stroke - Incidence in India and management of ischemic stroke. *Neurosciences.* 2002; 4(3):139-41.
7. P. Chitrabalam, DiptiBaskar, S. Revathy. A study on stroke in young and elderly in Rajiv Gandhi government general hospital, Chennai. *Int J Clin Med.* 2012;3:184-9.
8. Baiju Sam Jacob, Sreekumar B, V. Baby Paul. A clinical study of stroke in young in a teaching hospital. *J Evol Med Dent Sci.* 2014; 3(12):3199-204.
9. Bhattacharya S, Prasarsaha S, Basu A, Das K. A 5 year prospective study of incidence, morbidity and mortality stroke profile on stroke in a rural community of Eastern India. *J Indian Med Assoc.* 2005; 103(12):655-9.