

Harsukh Educational Charitable Society

International Journal of Community Health and Medical Research

Journal home page: www.ijchmr.com

doi: 10.21276/ijchmr

ISSN E: 2457-0117 ISSN P: 2581-5040

Index Copernicus ICV 2018=62.61

ORIGINAL RESEARCH

Assessment of dyslipidemia in type II diabetes mellitus patients

Siddharth Sonkar¹, Jyoti Pankaj²

¹Associate Professor, ²Assistant Professor, Department of General Medicine, Prasad Institute of Medical Science and Hospital Lucknow, U.P., India

ABSTRACT:

Background: Type II diabetes mellitus (T2DM), characterized by chronic hyperglycemia, impaired insulin secretion and insulin resistance. The objective of the present study was to study the lipid profile among type 2 diabetes mellitus patients. **Materials & Methods:** The present study was conducted in the department of general medicine. It comprised of 60 T2DM patients of both genders. Fasting blood sugar and lipid profile such as serum cholesterol, serum triglycerides, HDL, LDL and VLDL was assessed. **Results:** There were 30 males and females. The mean serum cholesterol level was 228.76 mg/dl, serum TG level was 202.6 mg/dl, HDL level was 39.8 mg/dl, LDL was 142.64 mg/dl and VLDL level was 43.5 mg/dl. **Conclusion:** The diabetes has a significant role in alteration of lipoprotein levels. There is significant alteration in lipid profile levels.

Key words: Diabetes, hyperglycemia, lipid.

Corresponding Author: Dr. Jyoti Pankaj, Assistant Professor, Department of General Medicine, Prasad Institute of Medical Science and Hospital Lucknow, U.P., India

This article may be cited as: Sonkar S, Pankaj J. Assessment of dyslipidemia in type II diabetes mellitus patients. HECS Int J Comm Health Med Res 2019; 5(2):79-81.

INTRODUCTION

Type II diabetes mellitus (T2DM), characterized by chronic hyperglycemia, impaired insulin secretion and insulin resistance, is caused by an interaction of genetic and environmental factors. T2DM is an important cause of mortality and morbidity worldwide and a major global public health problem.¹ The prevalence of T2DM has been increasing with the speeding up of urbanization and the improvement in living standards.² The number of persons with diabetes is expected to increase to 439 million by 2030, representing 7.7% of the total worldwide adult population 20–79 years old.³ Identifying patients at risk of T2DM is challenging, and we have no ideal therapy for T2DM or its chronic and serious complications, so prevention is of the utmost importance.

Type II diabetes mellitus (T2DM) has an intersecting underlying pathology with thyroid dysfunction. On one hand, thyroid hormones contribute to the regulation of carbohydrate metabolism and pancreatic function, and on the other hand, diabetes affects thyroid function tests to variable extents. Dyslipidaemia in individuals with type 2 diabetes is very common, with a prevalence of 72–85%.⁴ This phenomenon is associated with a significantly increased risk of coronary artery disease relative to

individuals without diabetes. Lipid abnormalities observed in patients with type II diabetes play a central role in the development of atherosclerosis. These lipid abnormalities are not only quantitative, but also qualitative and kinetic in nature.⁵ The objective of the present study was to study the lipid profile among type 2 diabetes mellitus patients.

MATERIALS & METHODS

The present study was conducted in the department of general medicine. It comprised of 60 Type II diabetes mellitus patients of both genders. The study was approved from institutional ethical committee. Patients were informed about the study procedure and written informed consent was taken.

A standard questionnaire including a detailed history of present and past medical conditions; family history of medical diseases; previous history of medications, alcohol, drug addiction and blood or blood product transfusion was taken. Fasting blood sugar and lipid profile such as serum cholesterol, serum triglycerides, HDL, LDL and VLDL was assessed. The data was collected systematically and analyzed statistically according to the standard statistical methods.

RESULTS

Table I Distribution of subjects

Gender	Male	Female
Number	30	30


Table I shows that there were 30 males and females.

Table II Assessment of lipid profile

Parameters (mg/dl)	Value
Serum cholesterol	228.76
Serum TG	202.6
HDL	39.8
LDL	142.64
VLDL	43.5

Table II, graph I shows that mean serum cholesterol level was 228.76 mg/dl, serum TG level was 202.6 mg/dl, HDI level was 39.8 mg/dl, LDL was 142.64 mg/dl and VLDL level was 43.5 mg/dl.

Graph I Assessment of lipid profile



DISCUSSION

Diabetes mellitus is one of the modern pandemics and an important health problem worldwide. Diabetes mellitus (DM) is a group of metabolic disease characterized by hyperglycemia resulting from defects in insulin secretion, insulin action or both. Uncontrolled diabetic patients are characterized by hyperglycemia, hyper insulinemia, protein glycation and oxidative stress which causes early appearance of diabetic complications. Type 2 diabetes mellitus (T2DM) is an important cause of mortality and morbidity worldwide and a major global public health problem. The prevalence of T2DM has been increasing with the speeding up of urbanization and the improvement in living standards.⁶ People with type 2 Diabetes have a high risk of cardiovascular diseases (CVD). Diabetic patients often exhibit an atherogenic

lipid profile, which greatly increases their CVD risk. However most of the individuals may also carry unnoticed dyslipidemia, characterized by increased levels of triglycerides and LDL and decreased HDL. The chronic hyperglycemia is associated with dysfunction, long-term damage and failure of various organs, especially the eyes, kidneys, nerves, heart and blood vessels.⁷ Population growth, urbanization, and increasing prevalence of obesity and physical inactivity are the major risk factors contributing to the increasing prevalence of type 2 diabetes (T2DM). Over the next two decades, the largest increase in the number of people with diabetes will be seen in developing countries, particularly in people of working age. Thyroid diseases and diabetes mellitus are the two most common endocrine disorders encountered in clinical practice which have been shown to mutually influence each other, and association between both the

conditions has long been reported.⁸ The present study was conducted to study the lipid profile among type 2 diabetes mellitus patients.

In present study both males and females had 30 patients each. The mean serum cholesterol level was 228.76 mg/dl, serum TG level was 202.6 mg/dl, HDL level was 39.8 mg/dl, LDL was 142.64 mg/dl and VLDL level was 43.5 mg/dl.

Subekti I et al⁹ in their study found that diabetes groups showed TC level of 316.18 mg/dl, TG of 358.36 mg/dl, LDL of 214.70 mg/dl, VLDL of 71.67 mg/dl and HDL of 29.80 mg/dl. LDL value was markedly high in diabetic HY patients. All the lipid profile parameters were significantly increased except HDL among the patients with diabetes and HY patients. Increase was more in cholesterol and LDL values among patients suffering from both diabetes and HY. HDL levels were lowest among the patients with diabetes and also decreased among the diabetic HYS.

CONCLUSION

The diabetes has a significant role in alteration of lipoprotein levels. There is significant alteration in lipid profile levels.

REFERENCES

1. Bali K, Vij AS. Pattern of dyslipidemia in type 2 diabetes mellitus in Punjab. *Int J Res Med Sci* 2016; 4:809-12.
2. Navneet Agrawal, Manoj Gulati. Study of Prevalence of

- dyslipidemia in Patients with Type 2 Diabetes Mellitus. *International Journal of Contemporary Medical Research* 2016; 8(3): 2212-14.
3. Zafar ME, Choudhary S, Rahman MF, Kumar R. Correlation of Dyslipidemia among Type II Diabetes Mellitus. *Sch. J. App. Med. Sci.*, 2016; 4(12A):4243-4248.
4. Jiffri EH. Relationship between lipid profile blood and thyroid hormones in patient with type 2 diabetes mellitus. *Adv Obes Weight Manag Control* 2017; 6(6):178-182.
5. Telwani AA, Wani ZH, Ashraf Y, Shah AA. Prevalence of dyslipidemia in type 2 diabetes mellitus: a case control study. *Int J Res Med Sci* 2017;5:4527-31.
6. Nirmala A. C., Edwin George, Rajendra Prasad. Comparison of lipid profile in type II diabetes mellitus patients with and without thyroid dysfunctions. *International Journal of Current Research* 2017; 9: 44952-44955.
7. Atin, Rav J, et al. Alteration in lipid profile among patients with type 2 diabetes mellitus. *Int J Res Med Sci* 2015; 27-35.
8. Ghosh A, Kundu D, Rahman F, Zafar ME, Prasad KR, Baruah HK, et al. Correlation of lipid profile among patients with hypothyroidism and type 2 diabetes mellitus. *BLDE Univ J Health Sci* 2018; 3:48-53.
9. Subekti I, Pramono LA, Dewiasty E, Harbuwono DS. Thyroid Dysfunction in Type 2 Diabetes Mellitus Patients. *Acta Medica Indonesiana*. 2018 Jan 19; 49(4):314.